



DOMINION[®] TREE & SHRUB

- Use on Lawns, Flowers, Ornamentals, Shrubs & Trees Including listed Fruit and Nut Trees
- One application kills white grubs, mole crickets and European crane fly larvae all season long
- Systemic insect protection for the entire plant



Apply at the tree base

NOT FOR SALE, USE OR DISTRIBUTION IN or INTO Nassau, Suffolk, Kings or Queens Counties, New York

For Outdoor Residential Use Only

| | |
|--------------------|---------|
| Active Ingredient | |
| Imidacloprid | 1.47% |
| Other Ingredients: | 98.53% |
| Total | 100.00% |

EPA Reg. No. 53883-205

EPA Est. No. 53883-TX-002

KEEP OUT OF REACH OF CHILDREN


CAUTION

See attached booklet for additional precautionary statements

NET CONTENTS: 32 FL. OZ. (1 QT.) (946 ML.)

Control Solutions, Inc.

5903 Genoa-Red Bluff • Pasadena, TX 77507-1041



Martin's[®]

Dominion[®] Tree & Shrub

For outdoor residential use only.

- Kills Insects & Prevents New Infestations
- Cannot Be Washed Off By Rain or Water
- Mix and Pour—Apply at the tree base

Active Ingredient:

Imidacloprid 1.47%

Other Ingredients: 98.53%

Total 100.00%

EPA Reg. No. 53883-205

EPA Est. No. 53883-TX-002

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See inside booklet for additional precautionary information.

Control Solutions, Inc.
5903 Genoa-Red Bluff • Pasadena, Texas 77507-1041

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

Note to Physician: No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to lakes, streams or ponds. Do not dump rinse water into sewers or other bodies of water. This pesticide is highly toxic to bees. Do not apply Dominion® Tree & Shrub when bees are active. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, be sure to read and follow all label directions. Shake well before using.

RESTRICTIONS:

- Read and follow directions when using.
- Do not spray the product into fish pools, ponds, streams, or lakes.
- Remove animal feeding dishes prior to application.

**ESTABLISHED ORNAMENTALS IN
RESIDENTIAL AREAS****Information**

Dominion® Tree & Shrub is for use on outdoor trees, shrubs and groundcovers in landscape plantings as listed in the table below:

| Landscape Plantings | Types of Plants |
|---|--|
| Roses, flowers, bedding plants and flowering shrubs | Azalea, begonia, camellia, cherry laurel, geranium, gardenia, hydrangea, impatiens, ligustrum, peonies, primrose, tea roses, salvia, zinnias and similar herbaceous plants |
| Evergreen and leafy shrubs and woody ornamentals | Arborvitae, boxwood, cotoneaster, euonymus, holly, juniper, laurel, privet, spruce yews and similar plants |
| Groundcovers | English ivy, Dichondra, ivy, pachysandra, vinca and other similar plants |

Dominion® Tree & Shrub is for use to kill the pests listed in the table below on outdoor trees and shrubs.

| Pests Controlled | |
|---|---|
| Adelgids | Leafhoppers (incl. Glassy-winged Sharpshooters) |
| Aphids | Leafminers (incl. Birch Leafminers) |
| Black Vine Weevil Larvae | Mealybugs |
| Roundheaded Borers (incl. Eucalyptus Longhorned Borers) | Pine Tip Moth Larvae |
| Flatheaded Borers (incl. Bronze Birch, Alder Borers) | Psyllids |
| Japanese Beetles (Adult) | Royal Palm Bugs |
| Lacebugs | Sawfly Larvae |
| Leaf Beetles (incl. Elm Leaf Beetles and Viburnum Leaf Beetles) | Scales (incl. Armored Scale [suppression] and Soft Scale) |
| | Thrips (suppression) |
| | Whiteflies |

Foliar Applications

Use Dominion® Tree & Shrub as a foliar spray to provide contact control of larvae, nymphs or adults of the following insect pests: adelgids, aphids, Japanese beetles, lace bugs, leaf beetles (including elm leaf beetles and viburnum leaf beetles), leafhoppers (including glassy-winged sharpshooter), mealybugs, psyllids, sawfly larvae, thrips (suppression) and whiteflies.

Application Instructions

Apply when insects first appear and before high pest populations are established. Retreat when re-infestation occurs, when high pest pressure is observed or repeat applications on an as-needed basis with 10-14 day intervals between applications.

How to Apply

Foliar applications with water must be made as a thorough cover spray to provide uniform distribution. Spray to wet all foliage (leaves, stems and branches). Target the undersides of the leaves and try to penetrate dense foliage with the spray droplets.

Hand-Operated Sprayers

Including backpack, compression, knapsack or tank type sprayers.

For foliar applications mix 1.5 (1½) teaspoons (tsp) of Dominion® Tree & Shrub per one (1) gallon of water.

| Spray Volume Gallons of Water | Teaspoons (tsp.) of product | Tablespoons (tbs.) of product |
|--|--|--|
| 1 gallon | 1½ tsp. | ½ tbs. |
| 2 gallons | 3 tsp. | 1 tbs. |
| 3 gallons | 4½ tsp. | 1½ tbs. |

1.0 fl. oz. = 2 tablespoons = 6 teaspoons (tsp.)

Hose-End Sprayer Instructions [For Gilmour Dial-A-Mix Sprayer or Air-O-Matic]

1. Set the dial to 1½ teaspoons (tsp.). Keep the dial set to this setting while spraying.
2. Add an adequate amount of product to the spray bottle to treat the total area. Do not add water to jar.
3. Spray to wet all foliage and direct the spray droplets for an even distribution on the target plants. Any unused product can be poured back into its original container.

Soil Applications

Soil application of Dominion® Tree & Shrub provides systemic activity and longer residual control as compared to foliar applications. Dominion® Tree & Shrub may be used on bedding plants, flowers, trees, shrubs, groundcovers and woody ornamentals in landscape plantings as a soil surface spray. Individual trees and shrubs may be treated with a soil drench.

| Pests Controlled | |
|------------------------------|--------------------------|
| Adelgids | Leafhoppers |
| Aphids | Leafminers |
| Armored scales (suppression) | Mealybugs |
| Black Vine Weevil Larvae | Pine Tip Moth Larvae |
| Eucalyptus Longhorned Borers | Psyllids |
| Flatheaded Borers | Royal Palm Bugs |
| Japanese Beetles (adult) | Sawfly Larvae |
| Lacebugs | Soft Scale (suppression) |
| Elm Leaf Beetles | Thrips (suppression) |
| Viburnum Leaf Beetles | White grub larvae |
| | Whiteflies |

Application Instructions

Seedbed or planting beds: Apply as a broadcast treatment to the soil and incorporate into the soil by cultivation or irrigation before seeding, planting and transplanting.

Established plantings: Apply Dominion® Tree & Shrub to the root zone of the established plants and irrigate the treated areas before the insect activity is expected.

Irrigate the treated area (0.5 inch) thoroughly after the application of Dominion® Tree & Shrub.

Hand-Operated Sprayers

Including backpack, compression, knapsack or tank type sprayers.

For soil applications to landscape plantings, mix 3.0 fl. oz. of Dominion® Tree & Shrub per one (1) gallon of water for treating approximately 333 sq. ft. Additional spray preparation rates are given in the table below.

| Spray Volume Gallons of Water | Fluid Ounces (fl. oz.) of Dominion® Tree & Shrub | Tablespoons (tbs.) of product | Area to be treated sq. ft. |
|--|---|--|---|
| 1 gallon | 3.0 fl. oz. | 6 tbs. | 333 sq. ft. |
| 2 gallons | 6.0 fl. oz. | 12 tbs. | 666 sq. ft. |
| 3 gallons | 9.0 fl. oz. | 18 tbs. | 1,000 sq. ft. |

1.0 fl. oz. = 2 tablespoons

Hose-End Sprayer Instructions

1. Measure the total area to be sprayed
2. Set the dial to 3 oz.
3. Add an adequate amount of product to the spray bottle to treat the total area, or add 9 fl. oz. of product for every 1,000 sq. ft. For example, pour 18 fl. oz. of product into sprayer jar to treat an area of 2,000 sq. ft. Do not add water to jar.
4. Spray evenly over the measured area. Any unused product can be poured back into its original container.

Soil Drench to Individual Trees and Shrubs

Dominion® Tree & Shrub quickly dissolves in water allowing the insecticide to move through the soil into the root zone. Once in the roots, the insecticide moves up through the tree or shrub providing year-long insect control. Dominion® Tree & Shrub will also provide control to new growth. Drench applications to individual trees will control aphids, borers, leafhoppers, lace bugs, mealybugs, scale insects, and other larvae as listed above.

For best results, use this product as part of a preventative treatment program before insect damage occurs. Use one soil drench treatment per year during early spring to early fall.

Dominion® Tree & Shrub can be applied anytime except when the ground is saturated or frozen. These conditions will make movement of the product down into the root zone difficult. When applying this product during dry periods, generously water the tree or shrub the night before treatment.

NOTE: For insect control, the insecticide must move throughout the tree or shrub. The time it takes to do this will depend on the size of the plant that is treated (one week to three months for very large trees). For this reason, soil drenches should occur 2-3 months before the expected pest infestation. Trees that are already heavily infested may not survive due to the existing pest damage and the resulting stress.

RESTRICTIONS:

Do not apply to soil that is frozen or waterlogged.

Keep children and pets off treated area until it has dried.

PREVENTATIVE TREATMENT INFORMATION AND INSTRUCTIONS FOR EMERALD ASH BORER AND ASIAN LONGHORNED BEETLE

When used as a soil injection or soil drench Dominion® Tree & Shrub **may** help to prevent infestation by Asian longhorned beetle and emerald ash borer. If being applied as a preventative measure in areas where these pests have been found, monitoring for pests must be conducted routinely. Follow use directions for applications to trees below.

Asian longhorned beetle and emerald ash borer are exotic pests subject to quarantine and eradication by USDA Plant Protection and Quarantine ("PPQ"). In the event an infestation is discovered, it must be reported to USDA PPQ (301-734-8717) or appropriate State or Local authorities, who will determine the necessary course of action.

Application Instructions

Soil Drench: Refer to the directions in the "TREES" section below.

Soil Injection: Apply the rates indicated in the table in the "TREES" section. Divide the appropriate volume by the number of holes required. Place holes at approximately evenly spaced intervals around the entire circumference of the tree at about 2/3 the distance from the trunk to the dripline. Inject at a depth of 8-12 inches.

TREES

Determining Amount of Dominion® Tree & Shrub to Use

Measure the circumference (distance around the tree trunk) in inches at chest height or at 4.5 feet from the soil using a flexible tape. Or, measure the diameter of the tree trunk in inches at chest height or at 4.5 feet from the soil.

Mix 1 oz. of Dominion® Tree & Shrub per inch of circumference of the tree trunk to one (1) gallon of water. Refer to the mixing chart below for the amounts of Dominion® Tree & Shrub needed.

How to Apply

Apply the solution to the root zone as a band around the base of the tree. Apply one (1) gallon of solution in a circular band from the base of the tree outward for two (2) feet. For trees with trunk diameters more than 16 inches at chest height, apply two (2) gallons of solution in a circular band from the base of the tree outward for three to four (3-4) feet.

Do not irrigate immediately after a soil drench application. Depending on the size and vigor of the tree, the uptake period may vary from one week to three months for larger trees. Insect control will be achieved when the uptake and translocation from the roots to all plant parts (leaves) has occurred.

| Tree Trunk Circumference (inches) | Tree Diameter (inches) | Amount of Dominion® Tree & Shrub (fl. oz.) | Volume of Water (gallons) | Band Width of Circular Area Around Base of Tree (outward feet) |
|-----------------------------------|------------------------|--|---------------------------|--|
| 8 | 2 | 8 | 1 | 2' |
| 10 | 3 | 10 | 1 | 2' |
| 15 | 5 | 15 | 1 | 2' |
| 20 | 6 | 20 | 1 | 2' |
| 25 | 8 | 25 | 1 | 2' |
| 30 | 10 | 30 | 1 | 2' |

NOTE: The Directions for Use, or this chart for 1 quart containers may not describe the spray preparation beyond 32 fl. oz. per the *EPA Label Review Manual* Chapter 11.

SHRUBS

Determining Amount of Dominion® Tree & Shrub to Use

Measure the height of the shrub to the nearest foot. Round any inches down to the nearest foot if less than 6" and up to the nearest foot if greater than 6".

Mix 3 oz. of Dominion® Tree & Shrub per foot of shrub height to one (1) gallon of water. Refer to table below for additional mixing instructions.

How to Apply

Apply the solution to the root zone as a band around the base of the shrub. Apply one (1) gallon of solution in a circular band from the base of the shrub outward for one (1) feet. For shrubs taller than 3 feet, apply one (1) gallon of solution in a circular band from the base of the shrub outward for two (2) feet.

Do not irrigate immediately after a soil drench application.

| Shrub Height (feet) | Amount of Dominion® Tree & Shrub (fl. oz.) | Volume of Water (gallons) | Band Width of Circular Area Around Base of Shrub (outward feet) |
|--------------------------------|---|--------------------------------------|--|
| 2 | 6 | 1 | 1' |
| 3 | 9 | 1 | 1' |
| 4 | 12 | 1 | 2' |
| 5 | 15 | 1 | 2' |
| 6 | 18 | 1 | 2' |
| 7 | 21 | 1 | 2' |
| 8 | 24 | 1 | 2' |

APPLE, PEAR, PEACH, PLUM, CHERRY AND PECAN TREES

For plants established in residential areas as ornamental plantings or as fruit/nuts for consumption.

| Plants | Insect Pests Controlled |
|--|---|
| Apple and Crabapple Pear, Common and Oriental | Aphids (except woolly apple aphid), Japanese beetles, leafhoppers (including glassy-winged sharpshooter), leafminers, mealybugs, and San Jose scale |
| Peach, Plum and Cherry | Aphids, cherry fruit fly maggot, Japanese beetles, leafhoppers, plant bugs, San Jose scale and stink bugs |
| Pecan | Yellow pecan aphid, black margined aphid, pecan leaf phylloxera, pecan spittlebug, pecan stem phylloxera |

Application Instructions

Timing of applications should coincide with vulnerable stages of the pests. The calendar schedules below are to be used as a guide to application timing but should not be considered a forecast or predictor for treating these insect pests. Consult your State Agricultural Extension Service agent for specific schedules and guidance for your local area. In addition, observe these guidelines for treating the following pests:

- **Aphids:** Make application when aphids first appear and before the leaves are rolled as a result of these pests.
- **Leafhoppers:** Make applications when leafhoppers first appear, during the nymph stage.
- **Leafminers:** Make two applications with intervals of 10 or more days. Make the first application soon after the petal fall is complete. A second application at the same dosage 10 days later should be applied to the early instar larvae.
- **Mealybugs:** Thorough and uniform coverage is necessary.
- **San Jose Scale:** Make application when crawlers are present. Ensure complete spray coverage of the branches, stems and leaves.
- **Spittlebug:** Apply when spittle masses are present and as new generations develop.

For Apples and Pears: Apply full cover sprays after the petal fall is complete. Continue application until the target insects are controlled. Apply up to five (5) foliar applications per season and allow 10 or more days between insecticide applications. Allow seven (7) days between the final application and the apple or pear harvest.

| Stage of Trees | Approximate Dates | Apple | Pear |
|---|-------------------|--------|--------|
| Petal fall | May 22 | | |
| 1st cover spray, 5-10 days after petal fall | June 1 | X | X |
| 2nd cover spray, 10-14 days later | June 15 | X | X |
| 3rd cover spray, 10-14 days later | June 29 | X | X |
| 4th cover spray, 10-14 days later | July 13 | X | X |
| 5th cover spray, 10-14 days later | July 27 | X | X |
| Waiting period before harvest | - | 7 days | 7 days |

For Peaches: Apply full cover sprays after the petal fall is complete. Continue application until the listed insects are controlled sufficiently. Use up to three (3) foliar applications per season. Allow at least 7 days between applications.

| Stage of Trees | Approximate Dates | Peach |
|---|-------------------|--------|
| Petal fall | May 25 – June 1 | |
| 1st cover spray, 5-10 days after petal fall | June 1 | X |
| 2nd cover spray, 7-14 days later | June 8 – 14 | X |
| 3rd cover spray, 7-14 days later | June 15 – 21 | X |
| Waiting period before harvest | - | 0 days |

For Cherries and Plums: Apply full cover sprays after the petal fall is complete. Continue application until the target insects are controlled sufficiently. Apply up to five (5) foliar applications per season and allow 10 or more days between insecticide applications. Allow seven (7) days between the final application and the cherry or plum harvest.

| Stage of Trees | Approximate Dates | Cherry | Plum |
|---|-------------------|--------|--------|
| Petal fall | May 22 | | |
| 1st cover spray, 5-10 days after petal fall | June 1 | X | X |
| 2nd cover spray, 10-14 days later | June 15 | X | X |
| 3rd cover spray, 10-14 days later | June 29 | X | X |
| 4th cover spray, 10-14 days later | July 13 | X | X |
| 5th cover spray, 10-14 days later | July 27 | X | X |
| Waiting period before harvest | - | 7 days | 7 days |

For Pecans: Apply full cover sprays after the petal fall is complete. Continue application until the listed insects are controlled sufficiently. Use up to three (3) foliar applications per season. Allow at least 7 days between applications.

| Stage of Trees | Approximate Dates | Pecan |
|---|-------------------|-------|
| Petal fall | Late April | |
| 1st cover spray, 5-10 days after petal fall | May 1 | X |
| 2nd cover spray, 10-14 days later | May 15 | X |
| 3rd cover spray, 10-14 days later | May 30 | X |

How to Apply

Foliar applications with water must be made as a thorough cover spray to provide uniform distribution. Spray to wet all foliage (leaves, stems and branches). Target the undersides of the leaves and try to penetrate dense foliage with the spray droplets.

Hose-End Sprayer Instructions

1. Set the dial to 1½ teaspoons (tsp.). Keep the dial set to this setting while spraying.
2. Add an adequate amount of product to the spray bottle to treat the infested trees. Do not add water to jar.
3. Spray to wet all foliage and direct the spray droplets for an even distribution on the target plants. Any unused product can be poured back into its original container.

Spray volume will vary with the size of the trees and thorough coverage is important. Small trees (less than 15 feet in height) will require approximately 2 gallons of diluted spray per tree. Larger trees (25 feet in height) will require approximately 8 gallons of diluted spray per tree.

Dilutions and Time Chart — The estimated amount of diluted spray and the time required to spray one tree with the dial setting of 1½ teaspoons (tsp.)

| Tree Height (feet) | Spray Volume per Infested Tree (gallons) | Estimated Time to Spray One Tree (minutes) |
|-------------------------------|---|---|
| 15 | 2 | 1 |
| 20 | 5 | 2½ |
| 25 | 8 | 4 |
| 30 | 15 | 7½ |
| 35 | 20 | 10 |

Soil Drench to Apple, Pear, Peach, Plum, Cherry and Pecan Trees

Dominion® Tree & Shrub quickly dissolves in water allowing the insecticide to move through the soil into the root zone. Once in the roots, the insecticide moves up through the tree or shrub providing year-long insect protection. Dominion® Tree & Shrub will also provide protection to new growth. Drench applications to individual trees will control aphids, borers, leafhoppers, lace bugs, mealybugs, scale insects, and other larvae as listed above.

For insect control, the insecticide must move throughout the tree or shrub. The time it takes to do this will depend on the size of the plant that is treated (one week to three months for very large trees). For this reason, apply soil drenches 2-3 months before the expected pest infestation.

For best results, use this product as part of a preventative treatment program before insect damage occurs. Use one soil drench treatment per year during early spring to early fall.

Dominion® Tree & Shrub can be applied anytime except when the ground is saturated or frozen. These conditions will make movement of the product down into the root zone difficult. When applying this product during dry periods, generously water the tree or shrub the night before treatment.

Do not apply to soil that is frozen or waterlogged.

Trees that are already heavily infested may not survive due to the existing pest damage and the resulting stress.

Determining Amount of Dominion® Tree & Shrub to Use

Measure the circumference (distance around the tree trunk) in inches at chest height or at 4.5 feet from the soil using a flexible tape. Or, measure the diameter of the tree trunk in inches at chest height or at 4.5 feet from the soil.

Mix 1 oz. of Dominion® Tree & Shrub per inch of circumference of the tree trunk to one (1) gallon of water. Refer to the mixing chart below for the amounts of Dominion® Tree & Shrub needed.

How to Apply

Apply the solution to the root zone as a band around the base of the tree. Apply one (1) gallon of solution in a circular band from the base of the tree outward for two (2) feet. For trees with trunk diameters more than 16 inches at chest height, apply two (2) gallons of solution in a circular band from the base of the tree outward for three to four (3-4) feet.

Do not irrigate immediately after a soil drench application. Depending on the size and vigor of the tree, the uptake period may vary from one week to three months for larger trees.

| Tree Trunk Circumference (inches) | Tree Diameter (inches) | Amount of Dominion® Tree & Shrub (fl. oz.) | Volume of Water (gallons) | Band Width of Circular Area Around Base of Tree (outward feet) |
|--|-------------------------------|---|----------------------------------|---|
| 8 | 2 | 8 | 1 | 2' |
| 10 | 3 | 10 | 1 | 2' |
| 15 | 5 | 15 | 1 | 2' |
| 20 | 6 | 20 | 1 | 2' |
| 25 | 8 | 25 | 1 | 2' |
| 30 | 10 | 30 | 1 | 2' |

NOTE: The Directions for Use, or this chart for 1 quart containers may not describe the spray preparation beyond 32 fl. oz. per the *EPA Label Review Manual* Chapter 11.

LAWNS

Information

Dominion® Tree & Shrub kills white grubs, mole crickets, European crane fly larvae and other pests listed in the table below. Dominion® Tree & Shrub forms a protective barrier in the soil which provides control of listed pests all season long. When applied early, Dominion® Tree & Shrub kills grubs, mole crickets and European crane fly before they hatch and damage lawns. Contact your County Extension Agent for the best dates to apply in your area. Dominion® Tree & Shrub may be sprayed over soil or mulch. Remove weed barrier before applying.

Use on lawns and yards including the following grasses:

| | |
|---|---|
| Kentucky bluegrass, Perennial ryegrass, Tall fescue, fescues (red or fine leaf) | Bermudagrass, Common and hybrid Centipedegrass St. Augustinegrass Zoysiagrass Buffalograss Bahigrass |
|---|---|

Pests Controlled

| Pest | Indication of Damage to Lawns |
|---|---|
| <p>Adelgids Annual bluegrass Weevil Aphids Billbugs Bluegrass Weevils Chinch bug (suppression) Cutworms (suppression) Elm leaf beetle European crane fly larvae Lacebugs Leafhoppers Leafminers Mealybugs Mole crickets Sawfly larvae Sod webworm Thrips (suppression) Whiteflies White Grubs (including larvae of Asiatic Garden Beetle, Black Turfgrass Ataenuis, Chafer, Japanese Beetle, May or June Beetle, and Oriental Beetle)</p> | <p>European crane fly larvae often cause damage by feeding on grass roots and disrupting the soil surface.</p> <p>Mole crickets often cause extensive damage by tunneling through lawns. Some species also feed on roots. Tunneling uproots the grass so it easily dries out. Lawns turn yellow, become thin, weedy and can eventually die.</p> <p>White grubs cause extensive damage to lawns by feeding on grass roots. Grass turns yellow, then brown and dies. Grubs chew off roots so that the grass is easily pulled up in pieces in your hands. You can roll it back like a carpet to reveal the grubs. If patches of dead grass appeared in your lawn late last summer, check for grubs.</p> |

Application Instructions

Treat for grubs from the beginning of May to the end of October, or when first noticed.

Contact your State Agricultural Extension Service agents for additional recommendations for your area.

RESTRICTIONS: Do not exceed one application of Dominion® Tree & Shrub per season for lawns.

RESTRICTIONS:

- Children and pets may re-enter the treated area after the spray has dried.
- Do not water lawn within one hour after applying.
- Do not apply Dominion® Tree & Shrub to waterlogged areas
- For grub (larvae) control, ½ inch of rainfall or irrigations should occur within 24 hours after application. Thoroughly wet the lawn a few hours after application.
- For armyworm and cutworm control, do not irrigate treated areas within 24 hours after application of this product.
- Do not mow treated lawn until after adequate rainfall or irrigation has occurred.

Hand-Operated Sprayers

Including backpack, compression, knapsack or tank type sprayers and low pressure boom sprayers.

For lawns, mix 2 to 3 fl. oz. of Dominion® Tree & Shrub per one (1.0) gallon of water for treating approximately 333 sq. ft. of lawn. Use the higher rate for heavier infestations of insect pests.

| Gallons of Water | Fluid ounces (fl. oz.) of Dominion® Tree & Shrub | Tablespoons (tbs.) of product | Area to be treated (sq. ft.) |
|------------------|--|-------------------------------|------------------------------|
| 1 gallon | 2.0 – 3.0 fl. oz. | 4 – 6 tbs. | 333 sq. ft. |
| 2 gallons | 4.0 – 6.0 fl. oz. | 8 – 12 tbs. | 666 sq. ft. |
| 3 gallons | 6.0 – 9.0 fl. oz. | 12 – 18 tbs. | 1,000 sq. ft. |

1.0 fl. oz. = 2 tablespoons

STORAGE AND DISPOSAL

Pesticide Storage: Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

Pesticide Disposal: If empty: Nonrefillable container. Do not reuse or refill this container. Rinse the empty product container thoroughly and disperse the rinse water on lawn as part of the application. Offer for recycling if available. If recycling is not available: then dispose of container in sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

If partly filled: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please call your local solid waste agency. Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Control Solutions, Inc.'s control it is impossible for Control Solutions, Inc. to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CONTROL SOLUTIONS, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Control Solutions, Inc. disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CONTROL SOLUTIONS, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

Martin's[®]

Dominion[®] **Tree & Shrub**

***For outdoor residential
use only.***

- Kills Insects & Prevents New Infestations
- Cannot Be Washed Off By Rain or Water
- Mix and Pour—Apply at the tree base

Active Ingredient:

Imidacloprid 1.47%

Other Ingredients: 98.53%

Total 100.00%

EPA Reg. No. 53883-205

EPA Est. No. 53883-TX-002

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See inside booklet for additional precautionary information.

Control Solutions, Inc.

5903 Genoa-Red Bluff • Pasadena, Texas 77507-1041